

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

11-6001: Polyclonal Antibody to TAB3

Clonality: Polyclonal

Application: WB
Reactivity: Human
Gene: TAB3
Gene ID: 257397
Uniprot ID: Q8N5C8
Format: Purified

Alternative Name: TAB3,MAP3K7IP3
Isotype: Rabbit IgG

Immunogen Information: A partial length recombinant TAB3 protein (amino acids 1-350) was used as the immunogen

for this antibody.

Description

TAB3 (TGF-beta activated kinase 1/MAP3K7 binding protein 3) has CUE domain and RanBP2-type zinc finger (NZF) domains mediates binding to two consecutive 'Lys-63'-linked ubiquitins. It contains 712 amino acids. TAB3 is a mediator of MAP3K7 activation and it forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation of TNF or IL-1. This successive MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kappaB transcription factor and AP1 transcription factor. This protein shows interaction with TAB1, TAB2, MAP3K7, TRAF2 and TRAF6. It has 2 isoforms produced by alternative splicing. Isoform 2 may be an oncogenic factor. TAB3 involved in necrosis and tumors like disease.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein A Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 4-6 ̸µg/ml




Fig-1: Western blot analysis of TAB3. Anti-TAB3 antibody (11-6001) was used at 4 μ g/ml on HepG2 lysate.