

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-4023: Monoclonal Antibody to HMGB1 (Clone: ABM24C2)

Clonality: Monoclonal **Clone Name:** ABM24C2 Application: FACS.WB Reactivity: Human Gene: HMGB1 Gene ID: 3146 **Uniprot ID:** P09429 **Purified** Format: HMGB1,HMG1 **Alternative Name:** Isotype: Mouse IgG2b Kappa

Immunogen Information: A partial length recombinant HMGB1 protein (amino acids 1- 200) was used as the

immunogen for this antibody.

## **Description**

HMGB1 (High-mobility group protein B1) is a 25 kDa non-histone chromatin protein, secreted by immune cells through leaderless secretory pathway. HMGB1 supports transcription and also interacts with nucleosomes to loosen packed DNA and remodel the chromatin. The presence of HMGB1 in the nucleus depends on posttranslational modifications. When the protein is not acetylated, it stays in the nucleus, but hyperacetylation on lysine residues causes it to translocate into the cytosol. HMGB1 has the ability to bend DNA in order to facilitate binding of other proteins to the DNA. HMGB1 has one important role in V(D)J recombination where act as a cofactor of the RAG complex. HMGB1 stimulates a plethora of cellular responses like inflammation, attracting inflammatory cells, recruiting stem cells and promoting their proliferation. HMGB1 is highly expressed in human kidney, tonsils, adipose tissue, adrenal cortex and amniotic fluid.

## **Product Info**

**Amount :**  $25 \mu g / 100 \mu g$ 

**Purification:** Protein G Chromatography

**Content:** 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 2-4 µg/ml, FACS analysis: 0.5 µg/10^6 cells



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-1: Western blot analysis of HMGB1. Anti- HMGB1 antibody (Clone: ABM24C2) was used at 2  $\mu$ g/ml on HepG2 lysate.




Fig-2: Intracellular flow analysis of HMGB1 in PBMC (Lymphocytes) using 0.5  $\mu$ g/10^6 cells of HMGB1 antibody (Clone: ABM24C2). Green represents isotype control; red represents anti-HMGB1 antibody. Goat anti-mouse PE conjugate was used as secondary.