

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-3003: Monoclonal Antibody to TLR9 (Clone: ABM1C51)

Clonality: Monoclonal
Clone Name: ABM1C51
Application: IHC,FACS,WB
Reactivity: Mouse,Human

Gene : TLR9
Gene ID : 54106
Uniprot ID : Q9NR96
Format : Purified

Alternative Name: TLR9,UNQ5798/PRO19605

**Isotype:** Mouse IgG1 Kappa

Immunogen Information: A partial length recombinant TLR9 protein (amino acids 100-290) was used as the immunogen

for the antibody.

## **Description**

TLR9, a member of toll-like receptor family are central to the innate immunity by identifying pathogen associated molecular patterns (PAMPs). TLR9 identify unmethylated CpG dinucleotides present in bacterial DNA leading to NF-kB activation.

## **Product Info**

**Amount:** 25 μg / 100 μg

**Purification:** Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 2-4 μg/ml, FACS analysis: 0.5 μg/10<sup>6</sup> cells, Immunohistochemical analysis: 5 μg/ml





Fig-1: Immunohistochemical analysis of TLR9 in human stomach tissue using TLR9 antibody (Clone: ABM1C51) at 5  $\mu$ g/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



\* abeomics

Fig-2: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 2  $\mu$ g/ml on (1) Raji and (2) EL4 lysates.




Fig-3: Intracellular flow analysis of TLR9 in human PBMC (Lymphocytes) using 0.5  $\mu$ g/10^6 cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary.




Fig-4: Intracellular flow analysis of TLR9 in Raji cells using 0.5  $\mu$ g/10^6 cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Fig-5 : Immunohistochemical analysis of TLR9 in human Liver tissue using TLR9 antibody (Clone: ABM1C51) at 20 µg/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-6: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 4  $\mu g/ml$  on Daudi lysate.