

90-2203: Human IL-32 (Interleukin 32) Pre-Coated ELISA Kit

Application : ELISA

Description

This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti- Human IL-32 antibody was precoated into 96-well plates. Biotin conjugated anti-human IL-32 detection antibody was used. Standards, test samples and biotin conjugated detection antibody were added to the wells subsequently. Wash buffer was used to wash any non-specific binding. HRP conjugated Streptavidin was used as secondary antibody. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Human IL-32 amount of samples captured in the plate. Optical Density (O.D) can be read at absorbance 450nm in a microplate reader. Concentration of Human IL-32 can be calculated using the standard curve.

Kit Components

| Item | Specifications | Storage |
|---|----------------|-------------|
| 96 well Strip ELISA Plate | 8 X 12 well | 4°C/-20°C |
| Lyophilized Standard | 2 vials | 4°C /-20°C |
| Sample and Standard Dilution Buffer | 20 ml | 4°C |
| Biotinylated Detection Antibody for hIL-32 | 120 µl | 4°C/-20°C |
| Antibody Dilution Buffer | 10 ml | 4°C |
| HRP Conjugated Streptavidin (SABC) | 120 µl | 4°C in dark |
| SABC Dilution Buffer | 10 ml | 4°C |
| TMB Substrate | 10 ml | 4°C in dark |
| Stop Solution | 10 ml | 4°C |
| 25X Wash Buffer | 30 ml | 4°C |
| Plate Sealer | 5 pieces | 2 |
| Product Manual | 1 | |

Product Info

| Amount : | 1 × 96 Tests |
|---------------------|-------------------------------|
| Content : | 1 × 96 well Format (96 tests) |
| Storage condition : | Please refer to the Manual |

For Research Use Only. Not for use in diagnostic/therapeutics procedures.



Application Note

This immunoassay kit allows for the in vitro quantitative determination of Human IL-32 concentrations in serum, plasma and other biological fluids.

Detection Range: 15.6 - 1000 pg/ml

Sensitivity: < 9.4 pg/ml




Fig-1:Human IL-32 Standard Curve