

## 20-1104: Polyclonal antibody to CARD8 (Tucan)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IHC,IP
<b>Reactivity :</b>	Human
<b>Gene :</b>	CARD8
<b>Gene ID :</b>	22900
<b>Uniprot ID :</b>	Q9Y2G2
<b>Format :</b>	Sera
<b>Alternative Name :</b>	CARD8,KIAA0955,NDPP1
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Full-length recombinant protein of human TUCAN (CARD8) was used as immunogen for this antibody

### Description

This antibody recognizes TUCAN, a 431 amino acid protein. TUCAN (tumor up-regulated CARD-containing antagonist of caspase nine) also known as CARD8 is a CARD domain containing protein. Proteins containing a CARD (caspase-associated recruitment domain) domain are key regulators of cell death, cell survival and cytokine production. TUCAN is an anti-apoptotic CARD protein that can protect tumors from cell death stimuli, and is overexpressed in certain forms of cancer. TUCAN has been shown to inhibit caspase-9 activation by binding to the CARD region of pro-caspase-9, thereby suppressing the formation of the Apaf-1-caspase-9 apoptotic complex and apoptosis. Additionally, a TUCAN isoform has been described that blocks both caspase-8 and caspase-9 mediated apoptosis. However, in some tumors, TUCAN play a role in modulating NFkB transcription factor survival signaling pathways.

### Product Info

<b>Amount :</b>	50 µl
<b>Content :</b>	50 µl sera
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200




Fig:1 Western blot analysis of TUCAN/CARD8 in tumor cell lines using 20-1104 at 1:2000. Cell lysates were normalized for total protein content.




Fig:2 Formalin-fixed, paraffin-embedded tissue human colon microarray stained for TUCAN/CARD8 expression at 1:2000 using 20-1104. A-A2, successively higher magnifications of the microarray. A, low magnification overview. A1, differential TUCAN/CARD8 staining between tissue cores is observed. A2, Areas of intense TUCAN/CARD8 staining in a colon carcinoma tissue core. Hematoxylin eosin counterstain.




Fig:3 Formalin-fixed, paraffin-embedded tumor/normal adjacent tissue cores from a human colon tissue microarray stained for TUCAN/CARD8 expression at 1:2000 using 20-1104. A and B are tumor tissue cores. A1 and B1 are the matched normal adjacent cores from A and B, respectively. Hematoxylin-eosin counterstain.