

## 20-1096: Polyclonal antibody to TRAF-2

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IHC,IP
<b>Reactivity :</b>	Human
<b>Gene :</b>	TRAF2
<b>Gene ID :</b>	7186
<b>Uniprot ID :</b>	Q12933
<b>Format :</b>	Sera
<b>Alternative Name :</b>	TRAF2,TRAP3
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	A synthetic peptide of human TRAF-2 ( amino acids 6-31 VTTPGSLELLQPGFSKTLGKLEAK) was used as immunogen for this antibody

### Description

This antibody recognizes TRAF2, which is a member of TRAF (TNF receptor-associated factor) adapter proteins composed 501 amino acids. It Links TNFRs (TNF Receptors) to the SAPKs (Stress-Activated Protein Kinases) and p38s. TRAF2 can activate ASK1 (Apoptosis Signal-Regulating Kinase-1) in vivo and can interact in vivo with the amino- and carboxyl-terminal noncatalytic domains of the ASK1 polypeptide. Expression of the amino-terminal noncatalytic domain of ASK1 can inhibit TNF and TRAF2 activation of SAPK. TRAF2 is a potential mediator of CD40 signaling. In vitro, TRAF2 and TRAF3 bind to the CD40 Cytoplasmic Tail (CD40CT) with much higher affinity than TRAF5 and TRAF6 (TNF Receptor Associated Factor-6) and that TRAF2 and TRAF3 bind to different residues of the CD40CT. The TRAF2-binding site of the CD40CT is critical for NF-KappaB and SAPK activation, as well as the UP-regulation of the ICAM1 (Intercellular Adhesion Molecule-1).

### Product Info

<b>Amount :</b>	50 µl
<b>Content :</b>	50 µl sera
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200




Fig:1 Western blot analysis of TRAF2 in normal human tissues using 20-1096 at 1:2000. TRAF2 is observed at ~60-64 kDa. Additional bands of lower molecular weight were seen in some cases, and may represent TRAF2 degradation fragments




Fig:2 Immunohistochemistry of formalin-fixed, paraffin-embedded human melanoma tissue section stained for TRAF2 using 20-1096 at 1:2000. A1 is a higher magnification of A. Hematoxylin-eosin counterstain.




Fig:3 Immunohistochemistry of formalin-fixed, paraffin-embedded human colon carcinoma tissue sections stained for TRAF2 using 20-1096 at 1:2000. Hematoxylin-eosin counterstain.