

10-7553: Monoclonal Antibody to Inhibin alpha (Clone: ABM44F5)

Clonality :	Monoclonal
Clone Name :	ABM44F5
Application :	WB,IHC
Reactivity :	Human
Gene :	INHA
Gene ID :	3623
Uniprot ID :	P05111
Format :	Purified
Alternative Name :	INHA
Isotype :	Mouse IgG2b Kappa
Immunogen Information	A partial length recombinant Inhibin alpha protein (amino acids 196-366) was used as the immunogen for this antibody.

Description

Inhibin alpha (INHA) are dimeric glycoproteins that are structurally and functionally related to transforming growth factorbeta and are composed of 1 Alpha-subunit and 1 of 2 beta-subunits (betaA or betaB). Inhibin, first identified as a hormone regulating pituitary FSH secretion, later recognized to act as a tumor suppressor in the gonad and adrenal glands. INHA are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins play important roles in normal gonadal function, including regulation of proliferation, differentiation, and steroidogenesis of Leydig and Sertoli cells via paracrine and autocrine processes.

Product Info

Amount :	25 µg / 100 µg
Purification :	Protein G Chromatography
Content :	25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Western blot analysis: 0.5-1; Immunohistochemical analysis-5-10 ÃDµg/ml



Fig-1: Western blot analysis of Inhibin Alpha. Anti-Inhibin Alpha antibody (Clone: ABM44F5) was tested at 0.5 μg /ml on Recombinant and human Ovary lysates.



₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Fig-2: Immunohistochemical analysis of Inhibin Alpha in Human Testis tissue using Inhibin Alpha antibody (Clone: ABM44F5) at 5 μ g/ml.