

36-1165: Monoclonal Antibody to CD15 / FUT4 (Reed-Sternberg Cell Marker)(FUT4/815 + BRA-4F1)

Clonality :	Monoclonal
Clone Name :	FUT4/815 + BRA-4F1
Application :	IHC
Reactivity :	Human
Gene :	FUT4
Gene ID :	2526
Uniprot ID :	P22083
Format :	Purified
Alternative Name :	FUT4,ELFT,FCT3A
Isotype :	Mouse IgM, kappa
Immunogen Information :	Purified human neutrophils (FUT4/815); Myelomonocytic leukemia cells (BRA-4F1)

Description


CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cells and some epithelial cells. CD15 antibody is very useful in the identification of Hodgkin's disease. CD15 is occasionally expressed in large cell lymphomas of both B and T phenotypes which otherwise have a quite distinct histological appearance.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes)



Formalin-fixed, paraffin-embedded human Hodgkin's Lymphoma stained with CD15 Monoclonal Antibody (FUT4/815 + BRA-4F1).